

VIM AND FDM FOR SOLVING BOLTZMANN EQUATION OF SELF- GRAVITATING GAS-DUST SYSTEM

NOR AZIRAN BINTI AWANG

**A thesis submitted in partial fulfillment of the award of the degree of
Master of Science (Mathematics)**

**Department Of Mathematics
Faculty Of Science
Universiti Teknologi Malaysia**

January, 2013

Specially dedicated to my beloved family
and those people who have given consistent support and guide me.

ACKNOWLEDGEMENT

I would like to say Alhamdulillah and thank all people who have helped and inspired me during my study.

First and foremost, I would like to record my gratitude to my honorable supervisor, Assoc Prof Norma Alias for his supervision, advice, and guidance from the very early stage of this research as well as giving me extraordinary experiences throughout the thesis. The most needed, he provided me persistent encouragement and support in various ways. I am truly grateful for having such a wonderful supervisor.

I would like to extend my appreciation my family who always support me in my journey of education. They have given me so much comfort, care and love, either financially or spiritually, of which word could not express and will forever be remembered in my heart.

Lastly, I would like to thank a lot to my course mates and other friends who have provided their support and assistance to enable the completion of this research.

ABSTRACT

In this research, Boltzmann equation of Self-Gravitating Gas-Dust System is introduced. The model consists of dust component equations, gas dynamics equations and also Poisson equation. The mathematical model is used in order to visualize the movement of the gas and dust in environment. This research only focuses on sequential algorithm and solving the model in 2 Dimension (2-D). Numerical solutions obtained by using the Variational Iteration Method (VIM) and Finite Difference Method (FDM). The results are computed by using Borland C++ Builder, MATLAB 2008a and MAPLE 13. Then, the graphs are visualized by using Microsoft excel. The result is based on numerical analysis. Numerical analysis under consideration is pattern of the graph, gradient of the graph, time execution and computational complexity. The movement of the gas-dust showed differently with the different method. At the end, the results show that the VIM and FDM are successfully in visualizing the movement of the gas-dust in reality. Based on theoretically, VIM is in solving gas-dust problem compared to FDM.

ABSTRAK

Dalam kajian ini, persamaan Boltzmann telah diperkenalkan. Model ini terdiri daripada persamaan habuk, persamaan gas dan juga persamaan Poisson. Model matematik ini digunakan untuk melihat pergerakan habuk dan gas dalam persekitaran. Kajian ini hanya memberi tumpuan pada algoritma berjujukan dalam 2 dimensi (2-D). Penyelesaian berangka telah diperolehi melalui kaedah lelaran variasi dan kaedah beza terhingga. Penyelesaian ini telah dilakukan dengan menggunakan perisian seperti Borland C++ Builder, MATLAB 2008a dan MAPLE 13 Manakala, graf dilakarkan menggunakan microsof excel. Keputusan adalah berdasarkan analisis berangka. Analisis berangka yang di kaji adalah corak pergerakan graf, kecerungan graf, masa pelaksanaan dan pengiraan komplek. Pergerakan habuk dan gas menunjukkan perbezaan apabila menggunakan kaedah penyelesaian yang berbeza. Hasil kajian menunjukkan kaedah lelaran variasi dan kaedah beza terhingga telah berjaya dalam menggambarkan pergerakan-pergerakan habuk-gas di alam realiti. Berdasarkan teori, kaedah lelaran variasi adalah keadah yang berkesan untuk menyelesaikan masalah habuk dan gas berbanding kaedah beza terhingga.